

Optical and radio variability of the northern VHE gamma-ray emitting BL Lac objects


ELINA LINDFORS

TUORLA OBSERVATORY, UNIVERSITY OF
TURKU

T.HOVATTA, K. NILSSON, R. REINTHAL,
V. FALLAH RAMAZANI, W. MAX-MOERBECK,
A. C. S. READHEAD, J. L. RICHARDS,
A. BERDYUGIN, A. SILLANPÄÄ, L. TAKALO



Motivation

- The most numerous extragalactic VHE gamma-ray sources are high synchrotron peaking BL Lac objects, which are rather little studied in the radio bands (and even optical bands)
- It is rather common that we only look at the “snapshots” for this sources (no long term studies)
- One-zone SSC is extremely widely used to model the blazar spectral energy distributions and typically the radio data is ignored (with a statement that it originates from another region)




The data



- 32 TeV-detected Northern BL Lac objects
- 15 GHz radio monitoring data from the OVRO telescope (<http://www.astro.caltech.edu/ovroblazars/>)
- Optical R-band monitoring from Tuorla blazar monitoring (<http://users.utu.fi/kani/1m>)



Optical vs. radio variability



- Radio and optical fluxes correlated
- Optical has larger variability amplitudes than the radio
 - Modulation index = σ / mean





Common origin of variations




17 / 27 objects show significant peaks in correlation analysis

- Median lag 70 days, optical leading radio
- Radio variations in the “core”!




13 / 31 sources show common rising or declining trends



Common slow variability?




> 50% flux has
common origin



> 25% flux has
common origin

Nature of the slow variability

- Red noise with the highest frequency variations missing from the radio data? NO
- Radio core that we see with VLBA? YES




Blue: Core flux from MOJAVE
(VLBA 15GHz)

SED modeling considerations

In about half of the sample sources at least some of the optical and radio emission (median >27 %) originates in the same emission region

- Single-zone models are not adequate for these objects
- Radio cannot always be ignored in SED modeling



Aleksic et al. 2014, A&A, 567, 135

Summary

About half of the Northern TeV-detected BL Lacs show a connection between optical and radio emission

- At least 27% of the emission in a common emission region
- Single-zone SED models not adequate
- Radio cannot be ignored

Looking at the other lower energy properties of the sample of VHE gamma-ray emitting population

- Work in progress
- Poster 19: V. Fallah Ramazani et al.